Your Guide to Understanding Genetic Conditions

C8B gene

complement C8 beta chain

Normal Function

The *C8B* gene provides instructions for making one piece, the beta subunit, of a protein complex called complement component 8. To form this complex, the beta subunit interacts with another piece made up of the alpha subunit (produced from the *C8A* gene) and the gamma subunit (produced from the *C8G* gene), which are linked.

Complement component 8 aids in a part of the body's immune response known as the complement system. The complement system is a group of proteins that work together to destroy foreign invaders, trigger inflammation, and remove debris from cells and tissues. Complement component 8 combines with several other complement proteins to form the membrane attack complex (MAC), which inserts itself in the outer membrane of bacterial cells. This complex creates a hole (pore) in the membrane, which kills the bacterium. This part of the immune response appears to be especially important for fighting infection by bacteria in the *Neisseria* genus.

Health Conditions Related to Genetic Changes

complement component 8 deficiency


Several mutations in the *C8B* gene cause complement component 8 deficiency type II. This condition is an immune system disorder, known as an immunodeficiency, in which the immune system is not able to protect the body effectively from foreign invaders such as bacteria. People with complement component 8 deficiency have a significantly increased risk of developing recurrent infections, particularly by *Neisseria meningitidis*, which causes meningitis, a serious condition that involves inflammation of the membranes surrounding the brain and spinal cord.

C8B gene mutations involved in complement component 8 deficiency are most often of a type called a C to T transition, in which a DNA building block (nucleotide) called cytosine (C) is changed to the nucleotide thymine (T). Most commonly, this change occurs in a region of the gene called exon 9, but it can occur in other regions. These mutations alter the sequence of protein building blocks, resulting in an abnormally short protein that, if produced, is likely broken down quickly. Other types of mutations can occur in the C8B gene; they also lead to an abnormally short protein that is quickly broken down. The resulting shortage of this protein impairs formation of complement component 8. Deficiency of this component prevents formation of membrane attack complexes. Without this part of the immune response, affected individuals are prone to infection, particularly by Neisseria bacteria.

Chromosomal Location

Cytogenetic Location: 1p32.2, which is the short (p) arm of chromosome 1 at position 32.2

Molecular Location: base pairs 56,929,210 to 56,974,383 on chromosome 1 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- C82
- complement component 8 subunit beta
- complement component 8, beta polypeptide
- complement component C8 beta chain isoform 1 preproprotein
- complement component C8 beta chain isoform 2
- complement component C8 beta chain isoform 3

Additional Information & Resources

Educational Resources

- Immunobiology: The Immune System in Health and Disease (fifth edition, 2001): The Terminal Complement Proteins Polymerize to Form Pores in Membranes that can Kill Certain Pathogens https://www.ncbi.nlm.nih.gov/books/NBK27100/# A186
- Molecular Biology of the Cell (fourth edition, 2002): Complement Activation Targets
 Pathogens for Phagocytosis or Lysis
 https://www.ncbi.nlm.nih.gov/books/NBK26846/#_A4679_

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28C8B%5BTIAB%5D%29+OR +%28complement+component+8,+beta+polypeptide%5BTIAB%5D%29%29+OR +%28%28C82%5BTIAB%5D%29+OR+%28complement+component+8+subunit +beta%5BTIAB%5D%29+OR+%28complement+component+C8+beta+chain+isoform+1+preproprotein%5BTIAB%5D%29+OR+%28complement+component +C8+beta+chain+isoform+2%5BTIAB%5D%29+OR+%28complement+component +C8+beta+chain+isoform+3%5BTIAB%5D%29+OR+%28complement+component +C8+beta+chain+isoform+3%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english %5BIa%5D+AND+human%5Bmh%5D

OMIM

 COMPLEMENT COMPONENT 8, BETA SUBUNIT http://omim.org/entry/120960

Research Resources

- ClinVar https://www.ncbi.nlm.nih.gov/clinvar?term=C8B%5Bgene%5D
- HGNC Gene Family: Complement system http://www.genenames.org/cgi-bin/genefamilies/set/492
- HGNC Gene Symbol Report http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/ hgnc_data.php&hgnc_id=1353
- NCBI Gene https://www.ncbi.nlm.nih.gov/gene/732
- UniProt http://www.uniprot.org/uniprot/P07358

Sources for This Summary

- Bubeck D. The making of a macromolecular machine: assembly of the membrane attack complex. Biochemistry. 2014 Apr 1;53(12):1908-15. doi: 10.1021/bi500157z. Epub 2014 Mar 18. Review. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/24597946
- OMIM: COMPLEMENT COMPONENT 8, BETA SUBUNIT http://omim.org/entry/120960
- Kaufman KM, Snider JV, Spurr NK, Schwartz CE, Sodetz JM. Chromosomal assignment of genes encoding the alpha, beta, and gamma subunits of human complement protein C8: identification of a close physical linkage between the alpha and the beta loci. Genomics. 1989 Oct;5(3):475-80.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/2613233

- Kaufmann T, Hänsch G, Rittner C, Späth P, Tedesco F, Schneider PM. Genetic basis of human complement C8 beta deficiency. J Immunol. 1993 Jun 1;150(11):4943-7.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/8098723
- Merle NS, Noe R, Halbwachs-Mecarelli L, Fremeaux-Bacchi V, Roumenina LT. Complement System Part II: Role in Immunity. Front Immunol. 2015 May 26;6:257. doi: 10.3389/ fimmu.2015.00257. eCollection 2015. Review.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/26074922
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4443744/
- Rosa DD, Pasqualotto AC, de Quadros M, Prezzi SH. Deficiency of the eighth component of complement associated with recurrent meningococcal meningitis--case report and literature review. Braz J Infect Dis. 2004 Aug;8(4):328-30. Epub 2004 Nov 19. Review. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/15565265
- Saucedo L, Ackermann L, Platonov AE, Gewurz A, Rakita RM, Densen P. Delineation of additional genetic bases for C8 beta deficiency. Prevalence of null alleles and predominance of C-->T transition in their genesis. J Immunol. 1995 Nov 15;155(10):5022-8.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/7594510

Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/gene/C8B

Reviewed: December 2015 Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services